Sexual well-being and Pelvic floor health

THE UNIVERSITY OF TEXAS MDAnderson Cancer Center

Making Cancer History®


Emma Holliday, MD Associate Prof GI Radiation Oncology

lets talk about sex, baby

• the problem


• the conversation

• the solutions



how does rectal cancer treatment cause sexual problems?

the problem(s)-





RT to the vagina

RT to the penile nerves/blood vessels

RT to the prostate





vaginal narrowing/ shortening

vaginal dryness


painful intercourse

decreased desire

erectile dysfunction

ejaculatory problems

Difficulty with orgasm & sexual satisfaction



MD Anderson Cancer Center



RT to the vagina

RT to the penile nerves/blood vessels

RT to the prostate

RT to the pelvis



surgical damage to nerves/ blood vessels vaginal narrowing/ shortening

vaginal dryness

painful intercourse

decreased desire

erectile dysfunction

ejaculatory problems

Difficulty with orgasm & sexual satisfaction









RT to the vagina

RT to the penile nerves/blood vessels

RT to the prostate

RT to the pelvis

surgical damage to nerves/ blood vessels

chemo-induced _____

vaginal narrowing/ shortening

vaginal dryness

painful intercourse

decreased desire

erectile dysfunction

ejaculatory problems

Difficulty with orgasm & sexual satisfaction







RT to the vagina


RT to the penile nerves/blood vessels

RT to the prostate

RT to the pelvis

surgical damage to nerves/ blood vessels


chemo-induced neuropathy












the problem(s)- problems caused by treatment?



Wallington & Holliday PRO 2021

not much data-especially for young patients



the conversation-

how well are doctors preparing
 patients for these issues?
(hint, we could do a lot better)

we've got to do better

prior to surgery- only **~50%** of patients report receiving satisfactory preop counseling and **<10%** of women receive information about postop sexual dysfunction.

prior to radiation- radiation oncologists only mention the possibility of sexual toxicity to~40% of patients

<10% who develop sexual problems post-treatment are referred

most resources are heteronormative in language and content

barriers to productive conversations

A lack of adequate training for doctors

A low comfort level discussing the subject

The assumption that sexuality is only about intercourse

The misconception that treatments are not available

how to self-advocate

informed decision-making

effective communication with health care providers

"I've heard rectal cancer treatment can impact my sexuality. What are the possible side effects of this treatment and how common are they?"

connected strength

Florez, ASCO knowledge conquers cancer

how to self-advocate

informed decision-making

effective communication with health care providers

connected strength

"These activities are important to my sex life and overall quality of life. Are there ways to be proactive with treatment to keep this in mind?

how to self-advocate

informed decision-making

effective communication with health care providers

connected strength

"are there professional or peersupport resources for this issue?

the solutions-

how can we be proactive in treating sexual and pelvic floor problems after treatment?

MD Anderson Cancer Center

Vaginal sexual dysfunction


Prevention:

- Vaginal dilators during radiation can limit damage
- After treatment dilator use at least 3X weekly for at least 2 yrs

Treatment:

- Vaginal moisturizers as part of regular self care
- Prescription Estrogen- improves vaginal blood flow and lubrication
- Clitoral stimulation program also improves vaginal blood flow
- Pelvic Floor PT- can help vaginismus, guided dilator therapy
- Sexual modification devices ("Oh-nut")

Arzola, Rooney... Holliday PRO 2023 Koerber et al Tumori 2019 Berman et al Int J Impotence Research 1999





Penile sexual dysfunction

Prevention:

- Prophylactic PDE5-i? Prostate data suggests 50mg sildenafil daily 3 days prior to RT through 6mo after RT was associated w/ lower rates of ED up to 12 mo (but not 18-24).
- Screen for low T and supplement if medically appropriate

Treatment:

- Treatment with PDE5-inibitors are effective in ~50% of men w/ post-RT ED
- 2nd line treatments include injectable mediations and penile implant.
- Anal dilator therapy for those engaging in receptive anal intercourse

Regardless of anatomy...

- Sexual rehabilitation counseling
- Individual or couples sex therapy
- Body image counseling
- Explore sexual modification devices with or without a partner

• Sex after cancer treatment may be different, but that doesn't mean it can't be meaningful and satisfying.

Thank you!

Happy to continue the conversation: ebholliday@mdanderson.org @DrEmmaHolliday